



Original Article



Profile and Assessment of Potential Risk Factors of Neonatal Candidemia Cases in a Tertiary-Care Hospital in North India

Safiya Firoze¹, Fatima Khan*, Shariq Ahmed¹, Asfia Sultan¹, Anees Akhtar¹, Edison Winsent¹, Uzma Firdaus², Zeeshan Ahmed¹, Bhanu Chaudhary³

1. Department of Microbiology, Jawaharlal Nehru Medical College (JNMC), Aligarh, Uttar Pradesh, India.

2. Department of Paediatrics, Jawaharlal Nehru Medical College (JNMC), Aligarh, Uttar Pradesh, India.

3. MBBS, Jawaharlal Nehru Medical College (JNMC), Aligarh, Uttar Pradesh, India.

* **Corresponding author:** Fatima Khan, Department of Microbiology, Jawaharlal Nehru Medical College (JNMC), Aligarh, 202001, Uttar Pradesh, India.

Email: fatimasalmanshah@gmail.com

Copyright: © Author(s). This is an open-access article distributed under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License \(CC BY-NC-ND 4.0\)](#), which permits non-commercial use, sharing, and reproduction in any medium or format, provided the original author and source are credited, and no modifications or adaptations are made.

ABSTRACT

Introduction: *Candida* blood stream infections (BSIs) are a major contributing factor to neonatal sepsis and sepsis-related morbidities. Despite advances in newborn care, there remains a considerable morbidity, a trend towards neonatal candidemia, and occasional cases of antifungal resistance.

Aim: Our study aimed to identify the prevalent *Candida* species, antifungal susceptibility (AFS) patterns, and to ascertain risk factors associated with candida BSIs in neonates.

Materials and Methods: This was an eighteen-month retrospective study of candidemia in neonates. Positive aerobic BactAlert blood cultures having yeast were speciated (HiMedia Chrom-Agar and Vitek2®). AFS of isolates was performed using VITEK2® and disc diffusion.

Results: 303 (11.1%) out of 2748 positive blood cultures revealed *Candida* species, from which 84 (27.7%) were confirmed as newborn (0 to 28 days) BSI cases. *Candida tropicalis* (29.8%) was the most common isolate, followed by *Candida albicans* (26.2%) and *C. glabrata* (25%). AFS of these isolates revealed overall resistance to amphotericin B of 2.4% and to fluconazole of 13.1%. 60 isolates tested by Vitek 2 were sensitive to voriconazole; there was 5% caspofungin, 1.7% micafungin, and 1.7% flucytosine resistance. Risk factor assessment included NICU admission (60.7%), broad-spectrum antibiotic usage (71.4%), low birth weight (65.5%), prematurity (65.5%), ventilator support (44%) and mortality (23.8%).

Conclusion: The rise in resistance and neonatal candidemia emphasises the necessity of reevaluating the application of stringent infection control methods, suitable prophylactic antifungals, and a limited antibiotic-use strategy, among other preventative, as well as therapeutic approaches.

KEYWORDS: Candidemia; Neonates; Antifungal; Resistance; Risk factors

INTRODUCTION

Newborn survival has significantly increased as a result of advancements in neonatal care. Nonetheless, both bacterial and mycological systemic infections

with early (within 72 hours) and late (post 72 hours) onset still pose a serious risk and are a major contributor to neonatal morbidity¹. There is growing

Citation: Firoze S, Khan F, Ahmed S, et al. Profile and Assessment of Potential Risk Factors of Neonatal Candidemia Cases in a Tertiary-Care Hospital in North India. *JASPI*. 2026;4(1):30-37. DOI: [10.62541/jaspi106](#)



recognition of the role of *Candida* species in neonatal intensive care units (NICUs). It is responsible for 9 to 13% of neonatal blood stream infections (BSI) and is the third most frequent cause of late-onset sepsis in NICU patients².

Although *Candida albicans* has traditionally been the most frequently isolated species, non-albicans *Candida* (NAC), including *C. glabrata*, *C. krusei*, *C. parapsilosis*, and *C. tropicalis*, have recently emerged as important opportunistic pathogens³. This shift in epidemiology is concerning, as NAC resistance to azoles is increasingly recognized and appears to be linked to the widespread use of azole agents⁴. Moreover, several NAC species demonstrate cross-resistance to newer triazoles, while others exhibit intrinsic resistance to fluconazole (FLZ). Hence, it is essential that all yeast isolates, whether from blood or other clinical specimens, undergo both antifungal susceptibility testing and species-level identification^{3,5}.

In neonatal intensive care units (NICUs), candidiasis is believed to arise primarily from endogenous colonization, given the high prevalence of fungal acquisition in this population. Within the first week of life, approximately 10% of neonates become colonized, a figure that may rise to nearly 64% after a four-week hospital stay^{6,7}. Outbreaks in NICUs have also been attributed to contaminated injectable preparations, such as lipid emulsions used for total parenteral nutrition, or via transmission from colonized healthcare personnel and patient-to-patient spread, particularly in the context of inadequate hand hygiene⁸⁻¹⁰. Other recognized risk factors include mechanical ventilation, broad-spectrum antibiotic therapy, prior gastrointestinal surgery, indwelling devices, low birth weight (LBW), and prematurity^{11,12}.

Clinically, *Candida* bloodstream infection often resembles bacterial sepsis, making diagnosis challenging. Mycotic sepsis may present with nonspecific features such as lethargy, glycemic instability, neutropenia, and respiratory distress. Disseminated fungal disease can result in extensive end-organ involvement, affecting the kidneys, liver, lungs, brain, eyes, bones, spleen, and joints. Notably, invasive infection may occur even in the absence of positive culture results¹³. Renal involvement may manifest as acute dysfunction, hypertension, or flank masses, while ocular dissemination can present as endophthalmitis, a severe complication requiring immediate medical intervention^{14,15}.

The crude death rate linked to candidemia is around 60%, whereas the attributable mortality rate may reach 49%. The at-risk group involved, the quality of medical care, and the resistance or

susceptibilities of different *Candida* species and their strains to certain antifungal drugs are some of the variables that can affect the incidence and related mortality of *Candida* BSI¹⁶. Owing to significant regional variations, local epidemiology expertise is essential for managing and preventing invasive infections caused by *Candida*¹⁷.

This study aimed to identify the various *Candida* pathogens, examine their sensitivity patterns, and assess risk factors associated with neonatal candidemia.

MATERIALS AND METHODS

Study Design and Setting: This (January 2023 to June 2024), retrospective-observational research was conducted at the Microbiology unit of a tertiary-care hospital located in a city of northern India. The study was approved by the Institutional Ethics Committee of Jawaharlal Nehru Medical College, Aligarh, India (IEC Approval No: IECJNMC/1078). *Candida* species were discovered in 303 (11.1%) of the 2748 blood culture bottles that were labelled positive during the study period (Figure 1). Of these, 90 were isolated from neonate's blood samples that ranged in age from 0 to 28 days. 6 samples were excluded because they lacked corroborated clinical characteristics or because they revealed yeast as a contamination or commensal. This research took into account 84 (27.7%) neonates who showed signs and symptoms of infection such as lethargy, feed intolerance, failure to thrive or sepsis combined with at least one positive blood culture for *Candida* species.

Laboratory Procedures: Blood specimens were obtained for culture, while adhering to aseptic procedures after obtaining informed consents from the guardians of the patients. Automated BacTAlert3D (Biomérieux, France) was used to perform the blood culture. A paired blood culture was taken. Upon flagging of the blood culture bottle, direct microscopy was done using Gram's stain followed by inoculation onto 5% sheep blood agar incubated at 37°C, along with two plates of Sabouraud dextrose agar incubated at 25°C and 37°C. Growth of colonies was visible in 24 hours for all cases. A number of phenotypic assays were run to determine the isolates of *Candida* species. These comprised of CHROMagar *Candida* (CHROMagar, Paris, France), chlamydospore formation on cornmeal agar (HiMedia, India), Gram's staining, and germ tube visualization.

Antifungal Susceptibility Testing: The Clinical and Laboratory Standards Institute (CLSI) standards¹⁸ were

followed in the disk diffusion method (Voriconazole - 1µg, Fluconazole - 25µg, Amphotericin B - 10µg, Caspofungin - 5µg, Itraconazole - 10 µg) used to determine the antifungal susceptibility of 24 isolates. Because the hospital serves a highly impoverished community, financial constraints prevented Vitek from being used on all samples. Vitek 2 Compact (Biomérieux, France) was used to identify 60 samples through the use of ID-YST and YS08 (Voriconazole - 0.12-8 µg/mL, Fluconazole - 1-64 µg/mL, Amphotericin B - 0.25-16 µg/mL and 5-Flucytosine - 1-64 µg/mL, Caspofungin - 0.125 -8 µg/mL, Micafungin - 0.06-8 µg/mL) cards for AFS testing.

Statistics: Statistical analysis was done using Statistical Package for Social Sciences (SPSS) version 20 and the prevalence of organisms was determined and expressed in percentage. Categorical variables were summarized with frequency and continuous variables were summarized with mean and standard deviation. For categorical outcomes, comparison between groups were performed using the Chi-square test or Fisher's exact test, as appropriate. A p-value of <0.05 was considered statistically significant.

OBSERVATIONS AND RESULTS

Neonates accounted for 90 of the 303 *Candida* isolates, of which 84 (27.7%) were proven to originate from cases of neonatal bloodstream infection. Likewise, among the 2445 aerobic bacterial isolates, 241 were from neonates, with 234 (9.6%) confirmed as being from neonate BSI cases (Figure 1 & Table 1).

Figure 1. Flagged Blood Culture Bottles and *Candida* in Blood Cultures from Neonates (0 to 28 days)




Table 1. Distribution of pure isolates from neonatal septicemia cases (n = 318)

	Organism	No. of isolates	Percentage
<u>Candida species</u> (n = 84)	<i>Candida albicans</i>	22	6.9%
	<i>C. glabrata</i>	21	6.6%
	<i>C. krusei</i>	4	1.3%
	<i>C. parapsilosis</i>	9	2.8%
	<i>C. pelliculosa</i>	3	0.9%
	<i>C. tropicalis</i>	25	7.9%
<u>Bacterial isolates</u> (n = 234)	<u>Enterobacteriaceae</u> (n = 84)		
	<i>Citrobacter sp.</i>	6	1.9%
	<i>Escherichia coli</i>	14	4.4%
	<i>Enterobacter sp.</i>	2	0.6%
	<i>Klebsiella</i>	59	18.6%
	<i>Serratia sp.</i>	3	0.9%
	<u>Gram positive cocci</u> (n = 106)		
	<i>Enterococci</i>	19	6%
	<i>Other Staphylococcus sp.</i>	62	19.5%
	<i>Staphylococcus aureus</i>	24	7.5%
	<i>Streptococci</i>	1	0.3%
	<u>Nil-fermenters</u> (n = 44)		
	<i>Acinetobacter</i>	21	6.6%
	<i>Pseudomonas</i>	21	6.6%
	<i>Stenotrophomonas sp.</i>	2	0.6%

The most common pathogens isolated were other *Staphylococcus* species (n= 62, 19.5%), *Klebsiella* species (n= 59, 18.6%), *Candida tropicalis* (n= 25, 7.9%), *Staphylococcus aureus* (n= 24, 7.5%), *C. albicans* (n= 22, 6.9%), and *C. glabrata*, *Acinetobacter* species, and *Pseudomonas* species (n= 21 each, 6.6%).

Non-albicans *Candida* species were responsible for 73.8% of the 84 newborn *Candida* BSI cases. *C. tropicalis* accounted for 25 (29.8%) of the total species (Table 2).

Table 2. Distribution of *Candida* species isolated from blood of neonates (n = 84)

Organism		No of isolates	Percentage
<i>Candida albicans</i>		22	26.2%
Non-albicans <i>Candida</i> (NAC)	<i>C. glabrata</i>	21	25%
	<i>C. krusei</i>	4	4.8%
	<i>C. parapsilosis</i>	9	10.7%
	<i>C. pelliculosa</i>	3	3.6%
	<i>C. tropicalis</i>	25	29.8%
		Total NAC = 73.8%	

The distribution of the other isolated species was: *Candida albicans*, 22 (26.2%); *C. glabrata*, 21 (25%); *C. parapsilosis*, 9 (10.7%); *C. krusei*, 4 (4.8%); and *Candida pelliculosa*, 3 (3.6%).

Table 3. Combined Antifungal Susceptibility Profile of *Candida*.

Organism	FLZ (n=84)		AMB (n=84)		VOR n=(60)		CSF n=(60)		MYC n=(60)		5FC n=(60)	
	S	R	S	R	S	R	S	R	S	R	S	R
<i>C. albicans</i> (22)	20/22 (90.9%)	2/19 (9.1%)	22/22 (100%)	0 (0)	19/19 (100%)	0 (0)	18/19 (94.7%)	1/19 (5.3%)	18/19 (94.7%)	1/19 (5.3%)	18/19 (94.7%)	1/19 (5.3%)
<i>C. glabrata</i> (21)	21/21 (100%)	0 (0)	21/21 (100%)	0 (0)	18/18 (100%)	0 (0)	18/18 (100%)	0 (0)	18/18 (100%)	0 (0)	18/18 (100%)	0 (0)
<i>C. krusei</i> (4)	0/4 (0)	4/4 (100%)	3/4 (75%)	1/4 (25%)	2/2 (100%)	0 (0)	2/2 (100%)	0 (0)	2/2 (100%)	0 (0)	2/2 (100%)	0 (0)
<i>C. parapsilosis</i> (9)	8/9 (88.8%)	1/9 (11.2%)	9/9 (100%)	0 (0)	5/5 (100%)	0 (0)	5/5 (100%)	0 (0)	5/5 (100%)	0 (0)	5/5 (100%)	0 (0)
<i>C. pelliculosa</i> (3)	3/3 (100%)	0 (0)	2/3 (66.7%)	1/3 (33.3%)	3/3 (100%)	0 (0)	3/3 (100%)	0 (0)	3/3 (100%)	0 (0)	3/3 (100%)	0 (0)
<i>C. tropicalis</i> (25)	21/25 (84%)	4/25 (16%)	13/13 (100%)	0 (0)	13/13 (100%)	0 (0)	11/13 (84.6%)	2/13 (15.4%)	13/13 (100%)	0 (0)	13/13 (100%)	0 (0)
	73 (86.9%)	11 (13.1%)	82 (97.6%)	2 (2.4%)	60 (100%)	0 (0)	57 (95%)	3 (5%)	59 (98.3%)	1 (1.7%)	59 (98.3%)	1 (1.7%)

Among the 24 *Candida* isolates that were tested for antifungal susceptibility by disk diffusion method (n=24), resistance to fluconazole, amphotericin B, was 20.8% and 0%, respectively. All of the *C. albicans*

isolates were sensitive to fluconazole and amphotericin B. No resistance was seen in *C. glabrata*, to fluconazole and amphotericin B. All of the 2 isolates of *C. krusei* were resistant to fluconazole (100%); no resistance was seen with amphotericin B. All the *C. parapsilosis* isolates were sensitive to all antifungals, except for 1 isolate showing FLZ resistance (25%). Out of 12 *C. tropicalis* isolates tested, resistance was seen with FLZ (2; 16.7%).

Among the 60 *Candida* isolates that were tested for antifungal susceptibility by Vitek 2 (Table 3), (n = 60), resistance to fluconazole (FLZ), amphotericin B (AMB), voriconazole (VOR), caspofungin (CSF), micafungin (MYC), and flucytosine (5FC), was 10%, 3.3%, 0%, 5%, 1.7%, and 1.7%, respectively. All of the *C. albicans* isolates were sensitive to amphotericin B and voriconazole. 2 (10.5%) isolates were resistant to FLZ, while 1 (5.3%) isolate of *C. albicans* was resistant to CSF, MYC, 5FC, each. All the isolates of *Candida glabrata* and *Candida parapsilosis* were sensitive to FLZ, AMB, VOR, CSF, MYC, and 5FC. Both isolates of *C. krusei* were resistant to fluconazole, and 1 (50%) was resistant to amphotericin B. No resistance was seen with VOR, CSF, MYC, and 5FC. Only 1 (33.3%) isolate of *C. pelliculosa* was resistant to amphotericin B, while 2 (15.4%) isolates of *C. tropicalis* were resistant to fluconazole and caspofungin, each. All 3 *C.*

pelliculosa isolates were sensitive to FLZ, VOR, CSF, MYC, and 5FC. All *C. tropicalis* isolates were sensitive to AMB, VOR, MYC, and 5FC.

Table 4. Demographic Data and Potential Risk Factors in Neonates

	Candidemia (n = 84)				Bacteraemia (n = 234)				P Value	
	No.		%		No.		%			
Admission										
NICU	51		60.7%		87		37.2%		<0.001	
Non-NICU	33		39.3%		147		62.8%		<0.001	
Antibiotics Usage										
< 7 days (d)	18	60	21.4%	71.4%	27	150	11.5%	64.1%	0.013*	
7 - 14 d	32		38.1%		40		17.1%		<0.001	
> 14 d	10		11.9%		83		35.5%		<0.001	
Not used	24		28.6%		84		35.9%		0.112	
Birth Weight										
< 1 kg [ELBW]	14	55	16.7%	65.5%	21	147	9%	62.8%	0.027*	
1 - 1.5 kg [VLBW]	16		19%		47		20.1%		0.419	
1.6 - 2.5 kg [LBW]	25		29.8%		79		33.8%		0.251	
> 2.5 kg	29		34.5%		87		37.2%		0.332	
Gender										
Males	52		61.9%		142		60.7%		0.422	
Females	32		38.1%		92		39.3%		0.422	
Gestational Age										
< 37 weeks (w)	55		65.5%		110		47%		0.002*	
≥ 37 w	29		34.5%		124		53%		0.002*	
Normal Vaginal Delivery	72		85.7%		210		89.7%		0.159	
Outcome										
Alive	64		76.2%		207		88.5%		0.003*	
Death	20		23.8%		27		11.5%		0.003*	
Ventilator Support										
< 7 d	7	37	8.3%	44%	57	104	24.4%	44.4%	0.001*	
7 -14 d	19		22.6%		31		13.2%		0.021*	
> 14 d	11		13.1%		16		6.8%		0.039*	
Not used	47		56%		130		55.6%		0.475	

ELBW= extremely low birth weight; VLBW= very low birth weight; LBW= low birth weight [* indicates P value <0.05]

A total of 318 neonates with bloodstream infection were analyzed, including 84 with candidemia and 234 with bacteremia (Table 4). NICU admission was significantly higher among neonates with candidemia compared to those with bacteremia (60.7% vs. 37.2%, $p<0.001$). Preterm birth (<37 weeks) was more frequent in the candidemia group (65.5% vs. 47%, $p=0.002$), with a significantly greater proportion of extremely low birth weight infants (<1 kg) (16.7% vs. 9%, $p=0.027$). Shorter prior antibiotic exposure (<7 days) was more common in candidemia ($p=0.013$),

whereas prolonged mechanical ventilation (>7 days) was significantly associated with candidemia ($p=0.021$). Mortality was significantly higher in the candidemia cohort compared with bacteremia (23.8% vs. 11.5%, $p=0.003$). No statistically significant differences were observed with respect to gender, mode of delivery, or normal birth weight categories.

DISCUSSION

Pathogens isolated in our positive neonatal BSI samples included *Staphylococcus* species (19.5%), *Klebsiella* species (18.6%), *Candida tropicalis* (7.9%), *Staphylococcus aureus* (7.5%), *Candida albicans* (6.9%), *Candida glabrata*, *Acinetobacter* species, and

Pseudomonas species (6.6% each). These findings closely parallel the observations of Nazir and Masoodi, who reported *Staphylococcus* species (16.6%), *Klebsiella* species (16.2%), *Candida tropicalis* (13.8%), *Acinetobacter* species (11.3%), *Staphylococcus aureus* (7.7%), and *Candida albicans* (5.6%) as the predominant isolates¹⁷. In the current research, 73.8% of the cases of newborn candidemia were caused by non-albicans *Candida* species, while 26.2% of cases were caused by *C. albicans*.

In our study, *Candida tropicalis* (29.8%) was the predominant species, followed by *C. albicans* (26.2%), *C. glabrata* (25%), *C. parapsilosis* (10.7%), *C. krusei* (4.8%), and *C. pelliculosa* (3.6%). Similar trends of *C. tropicalis* predominance have been reported nationally^{16, 19, 20}, including by Nazir and Masoodi¹⁷ and Basu et al.²¹, whereas Juyal et al. observed *C. parapsilosis* as most common³. Contrastingly, Tunc et al. in Turkey reported *C. albicans* predominance³⁰. The association of *C. tropicalis* with NICU outbreaks^{17, 23} underscores its virulence³. Fluconazole resistance (13.1%) was mainly among NAC, consistent with other Indian reports²⁴⁻²⁶, though all isolates remained 100% voriconazole susceptible¹⁹.

In contrast to our investigation, where fungal resistance to amphotericin B was found to be 2.4% across all *Candida* isolates, the Central India-based publication by Narain et al., showed that all fungal isolates were susceptible to the antifungals. Biswas et al., however, found comparable outcomes, citing a 3.6% resistance to amphotericin B¹⁹. Caspofungin, micafungin, and flucytosine had overall resistance of 5%, 1.7%, and 1.7%, respectively.

61.9 % were females and 38.2% were males, and 65.5% had low birth weight (<2.5 kg). Extremely low birth weight (≤ 1 kg) was significantly associated with candidemia (16.7%) compared to bacteremia (9%) ($p < 0.05$). Prematurity was also prominent, with 65.5% of candidemia cases occurring in infants born before 36 weeks versus 47% in bacteremia ($p < 0.05$). Notably, late preterm births (33-36 weeks) were significantly higher in candidemia (30.6%) than bacteremia (20.1%). Conversely, term births (>36 weeks) were significantly associated with bacterial BSIs (53% vs. 34.5%). These findings reaffirm preterm and low birth weight neonates as high-risk groups for candidemia^{12, 27}. In our study, prior antibiotic exposure was documented in 71.4% of neonates with candidemia. Importantly, antibiotic use for less than one week was statistically significant ($p < 0.05$) in candidemia (21.4%) compared with bacterial BSIs (11.5%). Our findings corroborate earlier studies by Fu et al.^{11, 12} and Benjamin et al.,

which demonstrated that broad-spectrum antibiotic exposure within seven days before diagnosis markedly increased candidemia risk in low-birth-weight neonates.

Ventilator support was required for 44% of the neonates with Candida BSI. Compared to neonatal bacteremia cases, which showed 13.2% of neonates requiring ventilatory assistance between 7 and 14 days, we found that 22.6% of neonatal candidemia patients required such ventilator support. This difference was statistically significant ($p < 0.05$). Furthermore, ventilator support for longer than two weeks was linked to 13.1% of occurrences of newborn candidemia. This finding was statistically significant ($p < 0.05$) in comparison to instances of newborn bacteremia, of which 6.8% required continuous ventilator support above 2 weeks.

In the present study, we observed several notable strengths in the form of identification of key risk factors such as low birth weight, prematurity, prior antibiotic exposure, and prolonged ventilator support adds important clinical relevance, further supported by statistical significance. However, certain limitations such as being a likely single-center observational study, the generalizability of the findings may be limited. Additionally, the absence of outcome measures such as mortality and lack of adjustment for potential confounders may influence interpretation.

CONCLUSIONS

This study underscores the significance of neonatal bloodstream infections, with a predominance of non-albicans *Candida* species, particularly *Candida tropicalis*, and the emergence of antifungal resistance patterns in NICU settings. Prematurity, low birth weight, prior antibiotic exposure, and prolonged ventilator support were identified as the key risk factors for candidemia.

These findings emphasize the need for strict infection control measures, including adherence to hand hygiene and aseptic practices, along with implementation of robust antimicrobial stewardship programs to limit unnecessary antibiotic use. Routine surveillance of pathogen distribution and antifungal susceptibility, early identification of high-risk neonates with consideration of targeted prophylaxis, strengthening of laboratory diagnostics, and regular staff training are recommended to improve neonatal outcomes.

ACKNOWLEDGEMENT

The technical help of Mr. Mohd Naved and Mr. Sanjay Sharma is duly acknowledged.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest

SOURCE OF FUNDING

None

ETHICS STATEMENT INCLUDING PATIENT CONSENT

Not applicable

AUTHORS' CONTRIBUTIONS

All authors contributed equally to the data collection and manuscript writing.

AI DECLARATION

No AI tool was used to generate research data, perform data analysis, interpret findings, formulate scientific conclusions, or make editorial decisions. All AI-assisted content was critically reviewed, verified, and revised by the authors. The authors take full responsibility for the accuracy, originality, integrity, and final content of the manuscript.

REFERENCES

1. Odabasi IO, Bulbul A. Neonatal Sepsis. Sisli Etfal Hastan Tip Bul. 2020 Jun 12;54(2):142–58. <https://doi.org/10.14744/semb.2020.00236>
2. Nirmal K, Das S, Jothisri L, Jain C, Singh N pal. P367 *Candida* sepsis in neonates a neglected pathogen: study from neonatal intensive care unit tertiary care hospital. Med Mycol. 2022 Sep 20;60(Suppl 1):myac072P367. <https://pmc.ncbi.nlm.nih.gov/articles/PMC9509727/>
3. Juyal D, Sharma M, Pal S, Rathaur VK, Sharma N. Emergence of Non-Albicans *Candida* Species in Neonatal Candidemia. N Am J Med Sci. 2013 Sep;5(9):541–5. <https://doi.org/10.4103/1947-2714.118919>
4. Lee Y, Puumala E, Robbins N, Cowen LE. Antifungal Drug Resistance: Molecular Mechanisms in *Candida albicans* and Beyond. Chem Rev. 2021 Mar 24;121(6):3390–411. <https://doi.org/10.1021/acs.chemrev.0c00199>
5. Berkow EL, Lockhart SR. Fluconazole resistance in *Candida* species: a current perspective. Infect Drug Resist. 2017 Jul 31;10:237–45. <https://doi.org/10.2147/idr.s118892>
6. Kelly MS, Benjamin DK, Smith PB. The Epidemiology and Diagnosis of Invasive Candidiasis Among Premature Infants. Clin Perinatol. 2015 Mar;42(1):105–17. <https://doi.org/10.1016/j.clp.2014.10.008>
7. Rao S, Ali U. Systemic fungal infections in neonates. J Postgrad Med. 2005;51 Suppl 1:S27-29. <https://pubmed.ncbi.nlm.nih.gov/16519252/>
8. Zingg W, Tomaske M, Martin M. Risk of Parenteral Nutrition in Neonates—An Overview. Nutrients. 2012 Oct;4(10):1490–503. <https://doi.org/10.3390/nu4101490>
9. Hamdan M, Puckett Y. Total Parenteral Nutrition. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 [cited 2023 Dec 5]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK559036/>
10. Omran EA, Eisa FF, Bakr WMK. Microbial Contamination of Neonatal Injectable Lipid Emulsions at 12 and 24 Hours' Infusion Time With Evaluation of Infection Control Measures. The Journal of Pediatric Pharmacology and Therapeutics: JPPT. 2020 Feb;25(1):53. <https://doi.org/10.5863/1551-6776-25.1.53>
11. Benjamin DK, Stoll BJ, Gantz MG, Walsh MC, Sanchez PJ, Das A, et al. Neonatal Candidiasis: Epidemiology, Risk Factors, and Clinical Judgment. Pediatrics. 2010 Oct;126(4):e865–73. <https://doi.org/10.1542/peds.2009-3412>
12. Fu J, Ding Y, Jiang Y, Mo S, Xu S, Qin P. Persistent candidemia in very low birth weight neonates: risk factors and clinical significance. BMC Infectious Diseases. 2018 Nov 12;18(1):558. <https://doi.org/10.1186/s12879-018-3487-9>
13. Delaloye J, Calandra T. Invasive candidiasis as a cause of sepsis in the critically ill patient. Virulence. 2014 Jan 1;5(1):161–9. <https://doi.org/10.4161/viru.26187>
14. Hope W, Natarajan P, Goodwin L. Invasive fungal infections. Clin Med (Lond). 2013 Oct;13(5):507–10. <https://doi.org/10.7861/clinmedicine.13-5-507>
15. Ly V, Sallam A. Fungal Endophthalmitis. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 [cited 2023 Dec 6]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK559257/>
16. Bhattacharjee P. Epidemiology and antifungal susceptibility of *Candida* species in a tertiary care hospital, Kolkata, India. Curr Med Mycol. 2016 Jun;2(2):20–7. <https://doi.org/10.18869/acadpub.cmm.2.2.5>
17. Nazir A, Masoodi T. Spectrum of *Candidal* species isolated from neonates admitted in an Intensive Care Unit of teaching hospital of Kashmir, North India. J Lab Physicians. 2018;10(3):255–9. https://doi.org/10.4103/jlp.jlp_1_18
18. Wayne PA. CLSI M44 Method for Antifungal Disk Diffusion Susceptibility Testing of Yeasts. 3rd ed.

- Pennsylvania, USA: Clinical and Laboratory Standards Institute; 2018. 44 p. chrome-extension://efaidnbmnnnibpcajpcglclefindmkaj/https://webstore.ansi.org/preview-pages/CLSI/preview_M44-A.pdf
19. Biswas B, Sharma AK, Seema K, Kumar A, Boipai M, Kumar M. Emerging threat of *Candida* resistance among neonates at a teaching institute of Jharkhand. *J Family Med Prim Care*. 2023 May;12(5):946–52. https://doi.org/10.4103/jfmprc.jfmprc_2104_22
 20. Tak V, Mathur P, Varghese P, Gunjiyal J, Xess I, Misra MC. The epidemiological profile of candidemia at an Indian trauma care center. *J Lab Physicians*. 2014 Jul;6(2):96–101. <https://doi.org/10.4103/0974-2727.141506>
 21. Basu S, Kumar R, Tilak R, Kumar A. *Candida* Blood Stream Infection in Neonates: Experience from A Tertiary Care Teaching Hospital of Central India. *Indian Pediatr*. 2017 Jul 15;54(7):556–9. <https://doi.org/10.1007/s13312-017-1067-5>
 22. Tunc G, Toksoz A, Kilicbay F. *Candidal* Infections in the Neonatal Intensive Care Unit: A Retrospective Observational Study. *Sisli Etfal Hastan Tip Bul*. 2023 Jun 20;57(2):204–9. <https://doi.org/10.14744/semb.2023.36037>
 23. Roilides E, Farmaki E, Evdoridou J, Francesconi A, Kasai M, Filioti J, et al. *Candida tropicalis* in a Neonatal Intensive Care Unit: Epidemiologic and Molecular Analysis of an Outbreak of Infection with an Uncommon Neonatal Pathogen. *J Clin Microbiol*. 2003 Feb;41(2):735–41. <https://doi.org/10.1128/jcm.41.2.735-741.2003>
 24. Xess I, Jain N, Hasan F, Mandal P, Banerjee U. Epidemiology of Candidemia in a Tertiary Care Centre of North India: 5-Year Study. *Infection*. 2007 Aug 1;35(4):256–9. <https://doi.org/10.1007/s15010-007-6144-6>
 25. Gupta N, Mittal N, Sood P, Kumar S, Kaur R, Mathur M. CANDIDEMIA IN NEONATAL INTENSIVE CARE UNIT. *Indian Journal of Pathology and Microbiology*. 2001 Jan;44(1):45. <https://pubmed.ncbi.nlm.nih.gov/12561995/>
 26. Kothari A, Sagar V. Epidemiology of *Candida* Bloodstream Infections in a Tertiary Care Institute in India. 2009 Oct 9; <https://doi.org/10.4103/0255-0857.49440>
 27. Hsieh E, Smith PB, Benjamin DK. Neonatal fungal infections: when to treat? *Early Hum Dev*. 2012 May;88(Suppl 2):S6–10. [https://doi.org/10.1016/s0378-3782\(12\)70004-x](https://doi.org/10.1016/s0378-3782(12)70004-x)