

Case Report

Extensive Deep Vein Thrombosis: A Rare Manifestation of Disseminated Tuberculosis

Shekhar Swaroop¹, Paras Singla², Upendra Baitha^{3*}, Gaurav Gupta³, Amandeep Singh³

1- Department of Gastroenterology, All India Institute of Medical Sciences (AIIMS), New Delhi, India.

2- Department of Paediatrics, All India Institute of Medical Sciences (AIIMS), New Delhi, India.

3- Department of Medicine, All India Institute of Medical Sciences (AIIMS), New Delhi, India.

* **Corresponding author:** Upendra Baitha, Department of Medicine, All India Institute of Medical Sciences (AIIMS), New Delhi, 110029, India.

Email: drupendrabaita@aiims.edu.in

Copyright: © Author(s). This is an open-access article distributed under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License \(CC BY-NC-ND 4.0\)](#), which permits non-commercial use, sharing, and reproduction in any medium or format, provided the original author and source are credited, and no modifications or adaptations are made.

ABSTRACT

A 20-year-old male with disseminated tuberculosis (pulmonary, pleural, and lymph node TB) presented with unusual manifestation in the form of extensive deep vein thrombosis (DVT) involving bilateral popliteal veins, femoral veins, iliac veins, and infra-hepatic inferior vena cava (IVC). Upon follow-up with the patient after two months of therapy, there was a significant clinical improvement, and edema and tenderness of the lower limb had decreased.

KEYWORDS: Tuberculosis; Diagnosis; Deep vein thrombosis

INTRODUCTION

It is well known that a chronic inflammatory state predisposes to deep vein thrombosis (DVT), but the literature on tuberculosis predisposing to DVT is scarce. Only a few case reports have reported the association between the two conditions. Tuberculosis is such a common infection in developing countries that clinicians should be aware that DVT can be one of the unusual manifestations of tuberculosis.

CASE REPORT

A 20-year-old male with no comorbidities or addictions presented with a history of dyspnoea, undocumented on and off low-grade fever, dry cough, significant weight loss and heaviness over the left side of the chest for the past five months, and painful swelling of bilateral lower limbs extending till thigh and visible veins over the abdomen for the past four months (Figure 1). There was no history of wheezing, bluish discoloration, palpitation, orthopnoea, or paroxysmal nocturnal dyspnoea.

On examination, the patient had tachypnoea and tachycardia. A general physical examination revealed bilateral enlarged posterior cervical lymph nodes, which were matted, tender, and firm in consistency without any redness or the local temperature rise over the lymph node, swelling, or discharging sinus or fistula. Examination of the lower limb revealed bilateral swollen and tender lower extremities extending to the thigh. Upon examination of the respiratory system, the trachea was shifted to the right side, with decreased expansion on the left side. On percussion, there was dullness over the left lung till the 4th intercostal space posteriorly and 5th intercostal space anteriorly. On auscultation, there was decreased air entry on the left side of the chest. Abdominal examination revealed dilated and engorged veins with a direction of blood flow from below upward, suggesting inferior vena cava obstruction.

On presentation, a chest X-ray was done, which revealed bilateral pleural effusion with massive

Citation: Swaroop S, Singla P, Baitha U, et al. Extensive Deep Vein Thrombosis: A Rare Manifestation of Disseminated Tuberculosis. *JASPI*. 2023;1(1):43-45. DOI: [10.62541/jaspi011](https://doi.org/10.62541/jaspi011)

left-sided pleural effusion. An intercostal drainage tube with multiple enlarged conglomerated necrotic supraclavicular, mediastinal, and periportal lymph nodes with bilateral empyema.

Figure 1: Collateral veins visible over patient's abdomen with extensive Deep Vein Thrombosis with disseminated Tuberculosis



Ultrasound screening was done for lower limb swelling, which revealed thrombosis of the bilateral femoral and popliteal vein, which was followed by a CT angiogram of the lower limb, which showed thrombotic occlusion of bilateral popliteal veins, femoral veins, iliac veins, and infra-hepatic IVC with collateral formation. Lymph node excisional biopsy revealed large areas of geographic necrosis with palisaded foamy histiocytes and few epithelioid cell granulomas. GeneXpert for MTB was positive and sensitive to rifampicin. We also considered other causes and risk factors for DVT, both acquired and inherited, and ruled them out with appropriate history, physical examination, and investigations.

The patient was started on anti-tubercular treatment (ATT) for disseminated tuberculosis and on low molecular weight heparin (LMWH) overlapped with warfarin for DVT and was monitored with international normalized ratio (INR) and the dose titrated accordingly. After starting ATT, the patient resolved symptoms and was discharged on the 15th day of ATT. LMWH and warfarin were stopped after eight days. On follow-up after two months of therapy, there was a decrease in lower limb swelling and tenderness, the fever had resolved, and the pleural effusion resolved.

DISCUSSION

Tuberculosis is one of the most common infections encountered in developing countries, and so is the case in India. DVT can have so many causes, including both inherited and acquired causes and inflammatory

diseases are one of them. Inflammatory diseases and infections have been known for a long to cause DVT.¹ Various mechanisms have been proposed for inflammation-induced thrombosis. Inflammation increases procoagulant factors and inhibits natural anticoagulant pathways and fibrinolytic activity, causing a thrombotic tendency. Chronic inflammation may cause endothelial damage, resulting in the loss of physiologic anticoagulant, antiaggregant, and vasodilatory properties of endothelium. On the other hand, coagulation also augments inflammation, mainly using thrombin-induced secretion of proinflammatory cytokines and growth factors. Platelets may also trigger inflammation by activating the dendritic cells.^{2,3}

However, the association of tuberculosis with DVT has begun to be reported in the last few years with only case series and case reports on the same.⁴⁻⁹ The mechanisms responsible for the development of DVT in patients with tuberculosis are not fully understood, and all three parts of Virchow's triad, including hypercoagulability, endothelial dysfunction, and stasis, may predispose to DVT in TB.⁶

Treatment of this condition requires special mention of the interaction of warfarin with rifampicin. CYP2C9 metabolizes warfarin, and rifampicin is an enzyme inducer. Co-administration of both warfarin and rifampicin can make it very difficult to reach the target INR with warfarin, and higher doses of warfarin may be required. Also, evidence is there to show that rifampicin itself can predispose to thrombosis risk.¹⁰ So, this aspect of treatment and whether we need to modify treatment for the same need to be studied. Our patient did not show any progress in thrombosis post-ATT. Also, INR was measured frequently, and with routine follow-up, warfarin was continued for three months without any complications.

CONCLUSIONS

Such an extensive thrombosis in a patient with disseminated tuberculosis without any retroperitoneal lymph node involvement and IVC obstruction has never been reported. Therefore, clinicians need to be aware that TB can predispose to extensive DVT and can be life-threatening because of the high risk of pulmonary embolism.

ACKNOWLEDGEMENT

None

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

SOURCE OF FUNDING

None

ETHICS STATEMENT INCLUDING PATIENT CONSENT

Not applicable

AUTHORS' CONTRIBUTIONS

SS: writing the draft

PS: review

UB: supervision

GG: editing

AS: conceptualization

AI DECLARATION

No AI tool was used to generate research data, perform data analysis, interpret findings, formulate scientific conclusions, or make editorial decisions. All AI-assisted content was critically reviewed, verified, and revised by the authors. The authors take full responsibility for the accuracy, originality, integrity, and final content of the manuscript.

REFERENCES

- Beristain-Covarrubias N, Perez-Toledo M, Thomas MR, Henderson IR, Watson SP, Cunningham AF. Understanding Infection-Induced Thrombosis: Lessons Learned From Animal Models. *Front Immunol.* 2019;10:2569. <https://pubmed.ncbi.nlm.nih.gov/31749809/>
- Branchford BR, Carpenter SL. The Role of Inflammation in Venous Thromboembolism. *Front Pediatr.* 2018;6:142. <https://pubmed.ncbi.nlm.nih.gov/29876337/>
- Aksu K, Donmez A, Keser G. Inflammation-induced thrombosis: mechanisms, disease associations and management. *Curr Pharm Des.* 2012;18(11):1478-93. <https://pubmed.ncbi.nlm.nih.gov/22364132/>
- Shitrit D, Fox L, Preiss R, Raz M, Mtvitzuk A. [Incidence of venous thromboembolism in 700 patients with acute tuberculosis]. *Harefuah.* 2012;151(4):208-54. <https://pubmed.ncbi.nlm.nih.gov/22616147/>
- Marjani M, Tabarsi P, Baghaei P, et al. Incidence of Thromboembolism in Hospitalized Patients With Tuberculosis and Associated Risk Factors. *Archives of Clinical Infectious Diseases.* 2012;7(2):56-9. <https://brieflands.com/journals/archcid/articles/20867>
- Gupta A, Mrigpuri P, Faye A, Bandyopadhyay D, Singla R. Pulmonary tuberculosis - An emerging risk factor for venous thromboembolism: A case series and review of literature. *Lung India.* 2017;34(1):65-9. <https://pubmed.ncbi.nlm.nih.gov/28144063/>
- Koç İ, Doğan S, Doğan Y, Ulasan A. [Co-existence of Lymph Node Tuberculosis and Pulmonary Embolism: A Case Report]. *J Clin Anal Med* 2015;6(suppl 5): 650-2. https://www.researchgate.net/publication/307844400_Co-existence_of_Lymph_Node_Tuberculosis_and_Pulmonary_EMBOLISM_A_Case_Report
- Kumar V, Gupta KB, Aggarwal R. Deep Vein Thrombosis in a Patient with Disseminated Tuberculosis. *J Infect Dis Antimicrob Agents.* 2011;28(1):63-7. https://scholar.google.com/citations?view_op=view_citation&hl=en&user=R3Mq1S8AAAAJ&citation_for_view=R3Mq1S8AAAAJ-f6ydRqryjwC
- Kouismi H, Laine M, Bourkadi J-E, Iraqi G. Association of deep venous thrombosis with pulmonary tuberculosis. *Egypt J Chest Dis Tuberc.* 2013;62(3):541-3. <https://www.sciencedirect.com/science/article/pii/S042276381300109X>
- White NW. Venous thrombosis and rifampicin. *Lancet.* 1989;2(8660):434-5. <https://pubmed.ncbi.nlm.nih.gov/2569610/>